Cow's milk allergy in infants is difficult to diagnose quickly and accurately.

In recent years, there have been concerns raised that cow's milk allergy is being over-diagnosed¹, it is therefore important to ensure that cow's milk allergy is identified accurately, and timely, appropriate management is put in place.

Example CMA symptoms

THERE ARE SEVERAL RESOURCES AND GUIDELINES available to healthcare professionals to ensure that the correct diagnosis is made.

Consider CMA in infants who exhibit 2 or more persistent and/or severe symptoms from the list $below^{1-3}$.





FOLLOW

THESE LINKS

to view validated

 Fox et al. Clin Transl Allergy. 2019;9:40 2. Luyt et al. Clin Experimental Allergy. 2014;642–672 3. Koletzko et al. JPGN. 2012;2:21-229 4. NICE Clinical Guideline 116 (2011). London; www.nice.org.uk/guidance/cg116 [09/03/20] 5. RCPCH (2011); London; www.rcpch.ac.uk/resources/allergy-care-pathway-food-allergy [09/03/20] 6. NICE Clinical Knowledge Summaries. Cow's milk allergy in children (2019); London;https:// cks.nice.org.uk/topics/cows-milk-allergy-in-children [09/03/20] 7. Lajeunesse et al. Arch Dis Chil Educ Pract Ed. 2019;104(6):286–291 8. Pulse Learning CPD for GPs. www.pulse-learning.co.uk/course/index. php?categoryid=260 [09/03/20]

20-067. Date of prep: March 2020. © Nutricia 2020.